

CENTRAL LIMIT THEOREMS FOR RANDOM POLYTOPES IN A SMOOTH CONVEX SET

VAN VU

ABSTRACT. Let K be a smooth convex set with volume one in $\mathbb{R}^d$. Choose n random points in K independently according to the uniform distribution. The convex hull of these points, denoted by K_n , is called a *random polytope*. We prove that several key functionals of K_n satisfy the central limit theorem as n tends to infinity.

1. INTRODUCTION

Let K be a convex set with volume one in $\mathbb{R}^d$. Choose n random points in K independently according to the uniform distribution. The convex hull of these points, denoted by K_n , is called a *random polytope*. The study of the key functionals (such as the volume, the number of vertices etc) of K_n , started by Efron [4] and Rényi and Sulanke [6], is a classical topic in convex geometry (e.g., see [13] for a survey).

Let $\text{Vol}(K_n)$ denote the volume of K_n . A well known conjecture in the field is that this random variable satisfies the central limit theorem as n tends to infinity.

Conjecture 1.1. *There is a function $\epsilon(n)$ tending to zero with n such that for every x*

$$\left| \mathbf{P}\left(\frac{\text{Vol}(K_n) - \mathbf{E}(\text{Vol}(K_n))}{\sqrt{\mathbf{Var}(\text{Vol}(K_n))}} \leq x\right) - \Phi(x) \right| \leq \epsilon(n),$$

where Φ denotes the distribution function of the normal distribution.

The conjecture has been verified only for the case when K is a ball in $\mathbb{R}^2$ [5].

In the study of random polytopes in a convex body K , the surface of K plays a critical role. A significant part of the literature focus on the following two cases:

- K is smooth, i.e., the boundary of K is twice differentiable with positive curvature bounded away from zero and infinity.

V. Vu is an A. Sloan Fellow and is supported by an NSF Career Grant.

- K is a polytope.

In this paper, we are going to prove the conjecture for the case when K is a smooth convex set.

Theorem 1.2. *Let K be a smooth convex body with volume one in $\mathbb{R}^d$. There is a function $\epsilon(n)$ tending to zero with n such that for every x*

$$\left| \mathbf{P}\left(\frac{\text{Vol}(K_n) - \mathbf{E}(\text{Vol}(K_n))}{\sqrt{\text{Var}(\text{Vol}(K_n))}} \leq x\right) - \Phi(x) \right| \leq \epsilon(n),$$

where Φ denotes the distribution function of the normal distribution.

Using the same method we can prove similar results for other functionals of K_n . Let $f_i(K_n)$ be the number of i -dimensional facets of K_n , where $0 \leq i \leq d-1$. For $i=0$, $f_0(K_n)$ is the number of vertices of K_n .

Theorem 1.3. *Let K be a smooth convex body with volume one in $\mathbb{R}^d$. There is a function $\epsilon(n)$ tending to zero with n such that for every x and every $0 \leq i \leq d-1$*

$$\left| \mathbf{P}\left(\frac{f_i(K_n) - \mathbf{E}(f_i(K_n))}{\sqrt{\text{Var}(f_i(K_n))}} \leq x\right) - \Phi(x) \right| \leq \epsilon(n),$$

where Φ denotes the distribution function of the normal distribution.

Remark 1.4. In both theorems one can take $\epsilon(n) = n^{-1/(d+1)+o(1)}$. See Remark 3.3.

Our approach is also applicable for the case when K is a polytope (pending on the availability of certain results). In fact, it can also be applied to other models of random polytopes; see Section 11) for a further discussion.

There are two main technical ingredients in our arguments. The first is recent results of Reitzner [7] which established central limit theorems for Poisson point processes. The second is (also recent) tail estimates by the author [11].

The paper is organized as follows. In the next section, we introduce the results from [7]. In Sections 3 and 4, we describe our general approach. In Section 5, we present the above mentioned tail estimates and a few corollaries. In Section 6, we prove a tail estimate on the difference $\text{Vol}(K_{n'}) - \text{Vol}(K_n)$, where n' and n are two large numbers which are relatively close to each other. As a corollary, we obtain a new estimate about $\mathbf{E}(\text{Vol}(K_{n'}) - \text{Vol}(K_n))$. The proof of Theorem 1.2 comes in Section 7. The proof of Theorem 1.3 appears in Sections 8 and 9. This latter proof is somewhat more complex due to a subtle geometrical fact (see the paragraph following Theorem 8.3). In Section 10, we summarize recent progresses on the distribution of $\text{Vol}(K_n)$ and $f_i(K_n)$. In Section 11, we discuss further applications of our method (for instance, the case when K is a polytope).

In the above sections we sometime omit technical details in order to maintain the flow of ideas. These details will be presented in the appendix at the end of the paper.

Notations. In the whole paper, we assume that n is large, whenever needed. The asymptotic notations are used under the assumption that $n \rightarrow \infty$. The hidden constants in O, Ω and Θ depend on the fixed convex body K . Vol and $\mathbf{P}$ denote volume and probability, respectively. Consider a (measurable) subset S of K and a random point x

$$\text{Vol}(S) = \mathbf{P}(x \in S).$$

$\mathbf{E}$ and $\mathbf{Var}$ denote expectation and variance, respectively; $\log$ denotes the logarithmic with natural base. For a point set P , $\text{Conv}(P)$ is the convex hull of P .

2. CENTRAL LIMIT THEOREMS FOR POISSON POINT PROCESSES

Let $X(n)$ be a Poisson point process in $\mathbb{R}^d$ of intensity n . Define the random polytope Π_n as the convex hull of the intersection of K with $X(n)$ (a precise definition is given in the next section). In a recent remarkable paper [7], Reitzner proved

Theorem 2.1. *Let K be a smooth convex body with volume one in $\mathbb{R}^d$. There is a function $\epsilon(n)$ tending to zero with n such that for every x*

$$\left| \mathbf{P}\left(\frac{\text{Vol}(\Pi_n) - \mathbf{E}(\text{Vol}(\Pi_n))}{\sqrt{\mathbf{Var}(\text{Vol}(\Pi_n))}} \leq x\right) - \Phi(x) \right| \leq \epsilon(n).$$

Theorem 2.2. *Let K be a smooth convex body with volume one in $\mathbb{R}^d$. Then for any $0 \leq i \leq d-1$ there is a function $\epsilon_i(n)$ tending to zero with n such that for every x*

$$\left| \mathbf{P}\left(\frac{f_i(\Pi_n) - \mathbf{E}(f_i(\Pi_n))}{\sqrt{\mathbf{Var}(f_i(\Pi_n))}} \leq x\right) - \Phi(x) \right| \leq \epsilon_i(n)$$

Remark 2.3. In Theorems 2.1 and 2.1, one can take

$$\epsilon(n) = cn^{-1/2+1/(d+1)} \log^{2+2/(d+1)} n$$

and

$$\epsilon_i(n) = cn^{-1/2+1/(d+1)} \log^{2+3i+2/(d+1)} n,$$

respectively [7].

In the same paper, Reitzner also determined the right order of magnitude of the variance of $\text{Vol}(K_n)$ and $f_i(K_n)$.

Theorem 2.4. *Let K be a smooth convex body with volume one in $\mathbb{R}^d$. Then $\mathbf{Var}(\text{Vol}(K_n)) = \Theta(n^{-(d+3)/(d+1)})$ and $\mathbf{Var}(f_i(K_n)) = \Theta(n^{(d-1)/(d+1)})$. The same estimates hold for Π_n .*

The upper bound for the variance was actually proved in an earlier paper [8]. A different proof was given by the current author in [11]. In [7], Reitzner proved a matching lower bound.

The proof in [11] also gives an upper bound for the moment of any fixed order. A matching lower bound follows from Theorems 1.2 and 1.3. For details see Section 10.

3. PASSING FROM THE POISSON MODEL TO THE UNIFORM MODEL

The heart of this paper is a technique which enables one to pass a statement for the Poisson model Π_n to a similar statement for the uniform model K_n . We will first focus on Theorem 1.2 concerning the volume.

In order to prove Theorem 1.2, given the results in the previous section, it suffices to show that $\text{Vol}(\Pi_n)$ and $\text{Vol}(K_n)$ have (approximately) the same expectation and variance. That is exactly what we are going to do

Proposition 3.1. *Let K be a smooth convex body with volume one in $\mathbb{R}^d$. Then there is a number $\delta = 1/(d+1) + o(1)$ such that the following holds.*

$$\mathbf{E}(\text{Vol}(\overline{\Pi}_n)) = \mathbf{E}(\text{Vol}(\overline{K}_n))(1 + O(n^{-\delta})),$$

where $\overline{\Pi}_n = K \setminus \Pi_n$ and $\overline{K}_n = K \setminus K_n$.

Proposition 3.2. *Let K be a smooth convex body with volume one in $\mathbb{R}^d$. Then there is a number $\delta = 1/(d+1) + o(1)$ such that the following holds.*

$$\mathbf{Var}(\text{Vol}(\Pi_n)) = \mathbf{Var}(\text{Vol}(K_n))(1 + O(n^{-\delta})).$$

Remark 3.3. Using the triangle inequality, Theorem 1.2 follows immediately from these propositions, Theorem 2.1 and the following lemma from [7]. Each of these results has an error term, but the error term $n^{-\delta}$ in the above proposition turns out to be the dominating one and it leads to Remark 1.4

Lemma 3.4. *Let K be a smooth convex body with volume one in $\mathbb{R}^d$. Then*

$$\left| \mathbf{P}(\text{Vol}(K_n) \leq x) - \mathbf{P}(\text{Vol}(\Pi_n) \leq x) \right| = O(n^{-2/(d+1)} \log^{2/(d+1)} n).$$

Remark 3.5. Proposition 3.1 has already been verified in [7]. Our proof here is different and works for every convex body K , without the smoothness assumption. It is mentioned in [7] that it has been conjectured that the variance of $\text{Vol}(\Pi_n)$ and

that of $\text{Vol}(K_n)$ are asymptotically the same, but no proof is known. Proposition 3.2 verifies this conjecture in a strong form.

In the following, we are going to focus on Proposition 3.2. Proposition 3.1 follows easily from the proof of Proposition 3.2.

Let us now give a better definition for the Poisson random polytope Π_n , mentioned in the previous section. In order to generate Π_n , one first generates a random number n' with respect to the Poisson distribution with mean n . Next, one generates a set P' of n' random points in K with respect to the uniform distribution. The convex hull of P' is Π_n .

It is well known from properties of the Poisson distribution that with high probability n' is close to n . To be precise,

$$\mathbf{P}(|n' - n| \geq A\sqrt{n \log n}) \leq n^{-A/4}$$

for any constant A (4 can be replaced by a better constant but this is not essential). Thus one may think about the distribution of Π_n as a distribution over $K_{n'}$, where $n' \in [n - A\sqrt{n \log n}, n + A\sqrt{n \log n}]$, for some large A .

Remark 3.6. We will prove rigorously that the contribution of those n' outside this interval is negligible. However, the reader can convince himself (herself) quickly on this point by observing that every quantity under investigation (such as the expectation or variance) is of order $\Omega(n^{-c})$ for some fixed c . Thus by setting A sufficiently large, omitting the n' outside the interval should not change our analysis significantly. We will use the phrase "with very high probability" to mean an event which holds with probability $1 - n^{-C}$ where we can set C arbitrarily large.

Our plan is to show that for any n' in the interval, the variance of $\text{Vol}(K_{n'})$ and that of $\text{Vol}(K_n)$ is very close to each other.

Lemma 3.7. *There is a number $\delta = 1/(d+1) + o(1)$ such that for any pair (n, n') where $n' = n + O(\sqrt{n \log n})$ and n is sufficiently large*

$$\mathbf{Var}(\text{Vol}(K_{n'})) = \mathbf{Var}(\text{Vol}(K_n))(1 + O(n^{-\delta})).$$

This, together with some simple arguments, implies Proposition 3.2.

4. ATTACKING LEMMA 3.7

We now describe the main idea of how to verify Lemma 3.7. This idea is quite general and works for other functionals (such as the number of vertices) as well.

Let us consider a number n' in the interval $[n - A\sqrt{n \log n}, A + \sqrt{n \log n}]$ for some large constant A . Assume (without loss of generality) that $n' > n$. Let Ω denote the product space K^n (equipped with the natural product measure). A point P in Ω is an ordered set $(x_1, \dots, x_n)$ of n random points (we generate the points one by one). The x_i are the coordinates of P . We use $Y(P)$ to denote the volume of the convex hull of P and μ to denote the expectation of $Y(P)$.

Remark 4.1. $Y(P)$ is, of course, just another way to express $\text{Vol}(K_n)$. It is however more convenient to use this notation in the proof below as it emphasizes the fact that Y is a function from Ω to $\mathbb{R}$.

Define Ω', P', μ' similarly (with respect to n'). The variance of $\text{Vol}(K_n)$ is

$$s = \int_{\Omega} |Y(P) - \mu|^2 \partial P$$

and the variance of $\text{Vol}(K_{n'})$ is

$$s' = \int_{\Omega'} |Y(P') - \mu'|^2 \partial P'.$$

Our goal is to show that δ

$$|s' - s| \leq n^{-\delta} s,$$

for some δ as claimed. We are going to use a coupling argument. Consider a point $P' = (x_1, \dots, x_{n'})$ in Ω' and the canonical decomposition

$$P' = P \cup Q$$

where $P = (x_1, \dots, x_n)$ and $Q = (x_{n+1}, \dots, x_{n'})$. In order to compare s and s' , we rewrite s as

$$s = \int_{\Omega'} (Y(P) - \mu)^2 \partial P'$$

where $Y(P)$ is understood as a function of P' which depends on the first n coordinates of P' . We now can write $s' - s$ as

$$s' - s = \int_{\Omega'} (Y(P') - \mu')^2 - (Y(P) - \mu)^2 \partial P'. \quad (1)$$

Next, we observe that

$$(Y(P') - \mu')^2 - (Y(P) - \mu)^2 = ((Y(P') - \mu') + (Y(P) - \mu))((Y(P') - \mu') - (Y(P) - \mu)).$$

Using a recent tail estimate in [11], we can prove that with very high probability, the quantities $|Y(P') - \mu'|$ and $|Y(P) - \mu|$ are small, of order $O(\sqrt{s} \log^{O(1)} n)$ in particular. This implies that typically

$$Y(P') - \mu' + (Y(P) - \mu) = O(\sqrt{s} \log^{O(1)} n) \quad (2)$$

and

$$(Y(P') - \mu') - (Y(P) - \mu) = O(\sqrt{s} \log^{O(1)} n). \quad (3)$$

These two inequalities, however, are not enough for our purpose. They only imply that the product of the left hand sides is of order $O(s \log^{O(1)} n)$, while we want something significantly small than s (the exponent in the logarithmic term is, unfortunately, positive). While the estimate for the sum $(Y(P') - \mu') + (Y(P) - \mu)$ seems to be near optimal, we can prove that a much better estimate holds for the difference $(Y(P') - \mu') - (Y(P) - \mu)$. In order to achieve this, we rewrite this difference as follows

$$(Y(P') - Y(P)) - (\mu' - \mu).$$

We are going to show that with very high probability

$$Y(P') - Y(P) = O(\sqrt{s} n^{-\delta}). \quad (4)$$

This will implies that $\mu' - \mu$ is of the same order. Thus, we gain a significant extra term $n^{-\delta}$ as claimed. In this part of the proof, the facts that P' contains P and $n' - n = O(\sqrt{n \log n})$ will play essential roles.

5. CONCENTRATION OF THE VOLUME

Again we use short hand Y to denote the random variable $\text{Vol}(K_n)$. Recall that the variance of Y is $\Theta(n^{-(d+3)/(d+1)})$. One expects that Y has sub-Gaussian tail, namely

$$\mathbf{P}(|Y - \mathbf{E}(Y)| \geq \sqrt{\lambda n^{-(d+3)/(d+1)}}) \leq \exp(-c\lambda)$$

for some constant c and a large range of λ . This intuition was confirmed in [11]. The following theorem is equivalent to Corollary 2.8 from that paper.

Theorem 5.1. *Let K be a smooth convex body with volume one in $\mathbb{R}^d$. There are positive constants c and α such that the following holds. For any $0 < \lambda \leq \alpha n^{\frac{(d-1)(d+3)}{(d+1)(3d+5)}}$*

$$\mathbf{P}(|Y - \mathbf{E}(Y)| \geq \sqrt{\lambda n^{-(d+3)/(d+1)}}) \leq 2\exp(-c\lambda) + \exp(-cn^{(d-1)/(3d+5)}). \quad (5)$$

It is useful to compare this theorem with the central limit theorem. In the central limit theorem, we have, for sufficiently large λ

$$\mathbf{P}(|Y - \mathbf{E}(Y)| \geq \sqrt{\lambda \mathbf{Var}(Y)}) \approx \frac{2}{\lambda} \exp(-\lambda/2). \quad (6)$$

Since $\mathbf{Var}(Y) = \Theta(n^{-(d+3)/(d+1)})$, the two bounds are comparable. The key difference here is that in the central limit theorem, λ has to be fixed and one takes n to infinity, while (5) also works for λ depending on n .

Remark 5.2. For the purpose of proving Theorem 1.2, it actually suffices to use a weaker theorem (Theorem 2.1 of [11]) whose proof is significantly simpler. In this weaker theorem instead of $\sqrt{\lambda n^{-(d+3)/(d+1)}}$ one has $\sqrt{\lambda n^{-(d+3)/(d+1)} \log n}$ (an extra logarithmic term). We state the above more precise theorem because we will discuss its analogue for $f_i(K_n)$ later.

We are also going to use the following standard fact, which can be routinely verified using the Laplace transform or properties of binomial coefficient (see [?], Chapter 2).

Lemma 5.3. *Let X be the sum of m i.i.d. indicator random variables. Then for any $t \geq 0$*

$$\mathbf{P}(X \geq \mathbf{E}(X) + t) \leq \exp\left(-\frac{t^2}{2(\mathbf{E}(X) + t/3)}\right).$$

6. DIFFERENCES OF VOLUMES

Consider two integers $n > n'$. We generate a set P' of n' random points as follows. First we generate a set P of n random points and next generate a set Q of $m = n - n'$ random points and then define $P' = P \cup Q$. Our goal is to show that if $n' - n$ is

small compared to n , then with very high probability, $Y(P') - Y(P)$ is small. The argument in this section works for every convex body K , *without* the smoothness assumption.

For a convex body K and a half space H , we call the intersection $H \cap K$ an ϵ -cap if $\text{Vol}(H \cap K) = \epsilon$. The union of all ϵ -caps is the ϵ -wet part of K . The complement of the ϵ -wet part is called the ϵ -floating body of K . We denote the ϵ -floating body and the ϵ -wet part by F_ϵ and $\overline{F}_\epsilon$, respectively. The volume of the ϵ -wet part play an important role and we denote it by ρ_ϵ .

Consider two points x and y in $\overline{F}_\epsilon$. We say that x sees y (with respect to F_ϵ) if the segment xy does not intersect the floating body F_ϵ . Let $S_{x,\epsilon}$ denote the set of those y that x sees. Define

$$g(\epsilon) = \sup_{x \in \overline{F}_\epsilon} \text{Vol}(S_{x,\epsilon}).$$

The set $S_{x,\epsilon}$ is the union of all ϵ -caps containing x . The quantity which is important to us is

$$\epsilon^* := \nu \log n / n$$

where ν is a sufficiently large constant.

Lemma 6.1. *Let K be a convex body of volume one in $\mathbb{R}^d$. For any constant β , there are constants γ and ν such that for $\epsilon^* = \nu \frac{\log n}{n}$, the following holds*

$$\mathbf{P}\left(Y(P') - Y(P) \geq (\gamma + 1)m\rho(\epsilon^*)g(\epsilon^*) + \gamma g(\epsilon^*) \log n\right) \leq n^{-\beta}.$$

Corollary 6.2. *Let K be a convex body of volume one in $\mathbb{R}^d$ and $\mu(n)$ the expectation of the volume of K_n . Then*

$$\mu(n + m) - \mu(n) = O(g(\epsilon^*)m\rho(\epsilon^*) + g(\epsilon^*) \log n). \quad (7)$$

If K is smooth, then

$$\mu(m + n) - \mu(n) = O((mn^{-2/(d+1)} + 1)n^{-1} \log^{O(1)} n). \quad (8)$$

If K is a polytope, then

$$\mu(m + n) - \mu(n) = O((mn^{-1} + 1)n^{-1} \log^{O(1)} n). \quad (9)$$

Remark 6.3. Using a stronger definition of smoothness, one can slightly improve (8). Indeed, if one assumes that $K \in \mathcal{K}_k^+$, then it is known that

$$\mu_n = \sum_{i=1}^k c_i n^{-(i+1)/(d+1)} + o(n^{-(k+1)/(d+1)})$$

where c_i are constants depending on K . Thus by taking k sufficiently large and using Taylor's expansion, one can prove (8) without the logarithmic term. Here $\mathcal{K}_k^+$ consists of convex body whose boundary is k time differentiable.

Proof In order to prove Lemma 6.1, we are going to need the following lemma from [2]. (For a different proof of this lemma using the VC dimension, see Section 4 of [11].)

Lemma 6.4. *There are positive constants c and c' such that the following holds for every sufficiently large n . For any $\epsilon \geq c' \log n/n$, the probability that K_n does not contain F_ϵ is at most $\exp(-c\epsilon n)$.*

Let E be the event that F_{ϵ^*} is not contained in $\text{Conv}(P)$. For any T

$$\mathbf{P}(Y(P') - Y(P) \geq T) \leq \mathbf{P}(E) + \mathbf{P}(Y(P') - Y(P) \geq T | \overline{E}).$$

By setting ν in the definition of ϵ^* we can assume that ϵ^* satisfies the condition of Lemma 6.4 and the probability $\exp(-c\epsilon^* n)$ we get from this lemma is at most $n^{-\beta-1}$. Thus we have

$$\mathbf{P}(Y(P') - Y(P) \geq T) \leq n^{-\beta-1} + \mathbf{P}(Y(P') - Y(P) \geq T | \overline{E}). \quad (10)$$

Observe that when $\overline{E}$ holds (namely, $F_{\epsilon^*} \subset \text{Conv}(P)$), the volume of $\text{Conv}(P)$ increases by at most $g(\epsilon^*)$ when we add to P a new point from $K \setminus \text{Conv}(P)$. Since $F_{\epsilon^*} \subset \text{Conv}(P)$, $K \setminus \text{Conv}(P) \subset \overline{F}_{\epsilon^*}$ and then

$$\mathbf{P}(Y(P') - Y(P) \geq T | \overline{E}) \leq \mathbf{P}(|Q \cap \overline{F}_{\epsilon^*}| \geq T/g(\epsilon^*)).$$

By substituting

$$T = g(\epsilon^*) \left((\gamma + 1)m\rho_{\epsilon^*} + \gamma \log n \right),$$

we have

$$\mathbf{P}(|Q \cap \overline{F}_{\epsilon^*}| \geq T/g(\epsilon^*)) = \mathbf{P}\left(|Q \cap \overline{F}_{\epsilon^*}| \geq (\gamma + 1)m\rho_{\epsilon^*} + \gamma \log n\right).$$

The quantity $|Q \cap \overline{F}_{\epsilon^*}|$ is a sum of m i.i.d. indicator random variables (each of which indicates the event that a random point belongs to the wet-part F_{ϵ^*} or not) with expectation

$$\text{Vol}(\overline{F}_{\epsilon^*}) = \rho_{\epsilon^*}.$$

Thus, the expectation of $|Q \cap \overline{F}_{\epsilon^*}|$ is $m\rho_{\epsilon^*}$. By applying Lemma 5.3 with

$$t = \gamma(m\rho_{\epsilon^*} + \log n),$$

we conclude that the probability in question is bounded from above by

$$\exp\left(-\frac{\gamma^2(m\rho_{\epsilon^*} + \log n)^2}{2(m\rho_{\epsilon^*} + \frac{1}{3}\gamma(m\rho_{\epsilon^*} + \log n))}\right).$$

It is simple to prove that for any $\gamma \geq 3$, the exponent

$$\frac{\gamma^2(m\rho_{\epsilon^*} + \log n)^2}{2(m\rho_{\epsilon^*} + \frac{1}{3}\gamma(m\rho_{\epsilon^*} + \log n))}$$

is bounded from above (with room to spare) by $\gamma/2$. Set γ large enough so that $\gamma/2 > \beta + 1$. We have

$$\exp\left(-\frac{\gamma^2(m\rho_{\epsilon^*} + \log n)^2}{2(m\rho_{\epsilon^*} + \frac{1}{3}\gamma(m\rho_{\epsilon^*} + \log n))}\right) \leq n^{-\gamma/2} \leq n^{-\beta-1}. \quad (11)$$

Since $2n^{-\beta-1} < n^{-\beta}$, (10) and (11) conclude the proof. $\blacksquare$

7. PROOF OF PROPOSITION 3.2

We first prove that there is a constant $\delta > 0$ such that for any n' where $|n' - n| = O(\sqrt{n \log n})$,

$$\mathbf{Var}(\text{Vol}(K_{n'})) = \mathbf{Var}(\text{Vol}(K_n))(1 + O(n^{-\delta})).$$

Without loss of generality, we can assume that $n' > n$. Let C be a large constant. We call a point $P' = (x_1, \dots, x_{n'}) \in \Omega'$ *C-typical* if the following hold

- $|Y(P') - \mu'| \leq C\sqrt{n^{-(d+3)/(d+1)} \log n}$

- $|Y(P) - \mu| \leq C\sqrt{n^{-(d+3)/(d+1)} \log n}$
- $Y(P') - Y(P) \leq Cg(\epsilon^*)(m\rho_{\epsilon^*} + \log n)$

The results in the previous two sections show that there is a positive constant α independent of C such that P' is C -typical with probability at least $1 - n^{-\alpha C}$. Since K is smooth, $g(\epsilon) = O(\epsilon)$ and $\rho(\epsilon) = \Theta(\epsilon^{2/(d+1)})$ for all sufficiently small ϵ . Substituting $\epsilon^* = \Theta(\log n/n)$ and $m = O(\sqrt{n \log n})$, we obtain

$$Cg(\epsilon^*)(m\rho_{\epsilon^*} + \log n) = O(n^{-(d+5)/2(d+1)} \log^{O(1)} n).$$

This implies that if P' is C -typical then

$$\begin{aligned} |(Y(P') - \mu')^2 - (Y(P) - \mu)^2| &= O\left(\sqrt{n^{-(d+3)/(d+1)} \log n} \times n^{-(d+5)/2(d+1)} \log^{O(1)} n\right) \\ &= O\left(n^{-(d+4)/(d+1)} \log^{O(1)} n\right). \end{aligned}$$

On the other hand, the variance s is of order $\Theta(n^{-(d+3)/(d+1)})$. Thus, we obtain the claim of Lemma 3.7 with $\delta = 1/(d+1) - o(1)$.

Now we are going to conclude the proof of Proposition 3.2. Consider the interval $I = [n - A\sqrt{n \log n}, n + A\sqrt{n \log n}]$. For every number n' in this interval, let $E_{n'}$ denote the events that n' is sampled (according to the Poisson distribution with mean n) and E_0 denote the event that the sampled number does not belong to the interval. We have

$$\mathbf{Var}(\Pi_n) = \mathbf{E}_{n'}(\mathbf{Var}(\text{Vol}(\Pi_n)|E_{n'})) + \mathbf{Var}(\mathbf{E}(\text{Vol}(\Pi_n)|E_{n'})),$$

where $n' \in I$ or $n' = 0$. By increasing A , we can assume that $\mathbf{P}(E_0) \leq n^{-B}$ for a sufficiently large B . Thus, the contribution of $\text{Vol}(\Pi_n|E_0)$ is negligible. Moreover, for any $n' \in I$,

$$\text{Vol}(\Pi_n)|E_{n'} = \text{Vol}(K_{n'}).$$

Thus, we have

$$\mathbf{Var}(\text{Vol}(\Pi_n)|E_{n'}) = \mathbf{Var}(\text{Vol}(K_{n'})) = \mathbf{Var}(\text{Vol}(K_n))(1 + O(n^{-\delta}))$$

and also

$$\mathbf{E}(\text{Vol}(\Pi_n)|E_{n'}) = \mathbf{E}(\text{Vol}(K_{n'})) = \mathbf{E}(\text{Vol}(K_n))(1 + n^{-\delta}).$$

These imply that

$$\mathbf{Var}(\text{Vol}(\Pi_n)) = \mathbf{Var}(\text{Vol}(K_n))(1 + O(n^{-\delta})),$$

proving Proposition 3.2. $\square$

8. PROOF OF THEOREM 1.3

We first prove this theorem for the case $i = 0$ ($f_0(K_n)$ counts the number of vertices of K_n). The (slightly more technical) proof for a general i follows in the next section. We need to prove the following proposition.

Proposition 8.1. *Let K be a smooth convex body with volume one in $\mathbb{R}^d$. Then there is a number $\delta = 1/(d+1) + o(1)$ such that the following holds.*

$$\mathbf{Var}(f_0(\Pi_n)) = \mathbf{Var}(f_0(K_n))(1 + O(n^{-\delta})).$$

Once the proposition is available, one can use Theorem 2.2 to conclude the proof. The above proposition follows from a variant of Lemma 3.7.

Lemma 8.2. *There is a number $\delta = 1/(d+1) + o(1)$ such that for any pair (n, n') where $n' = n + O(\sqrt{n \log n})$ and n is sufficiently large*

$$\mathbf{Var}(f_0(K_{n'})) = \mathbf{Var}(f_0(K_n))(1 + O(n^{-\delta})).$$

In order to prove this lemma, we use the ideas described in Section 4. We are going to use short hand Z to denote $f_0(K_n)$. Theorem 5.1 is replaced by the following result, which is a corollary of Theorem 2.11 from [11].

Theorem 8.3. *Let K be a smooth convex body with volume one in $\mathbb{R}^d$. There are positive constants c and α such that the following holds. For any $0 < \lambda < \alpha n^{\frac{d-1}{3(d+1)}}$, we have*

$$\mathbf{P}(|Z - \mathbf{E}(Z)| \geq \sqrt{\lambda n^{(d-1)/(d+1)}}) \leq 2 \exp(-c\lambda) + \exp(-cn^{(d-1)/(3d+5)}).$$

While it seems that we have all necessary tools in hand, the proof of Lemma 8.2 is still not a straightforward modification of the proof of Lemma 3.7. In fact this proof is much more delicate due to the following subtle obstruction: Z is not a *monotone* random variable. Recall that when we bound the difference between the volumes $Y(P')$ and $Y(P)$, we relied on the following observation: Adding a point from $P' \setminus P$ to P cannot increase $Y(P)$ by too much, unless P itself is very distorted, which happens with a negligible probability. When we repeat this argument for the number of vertices instead of the volume, we have to take into account the fact

that the number of vertices of $\text{Conv}(P)$ can actually *decrease* when we add a new point to P . Therefore, we have to give an upper bound for both $Z(P') - Z(P)$ and $Z(P) - Z(P')$.

The handling of $Z(P') - Z(P)$ is similar to that of $Y(P') - Y(P)$. If $\text{Conv}(P)$ contains the floating body F_{ϵ^*} , then a point from $Q = P' \setminus P$ would increase $Z(P)$ by at most one if it falls in the wet part $\overline{F}_{\epsilon^*}$. Thus, $Z(P') - Z(P)$ (conditioned on P contains F_{ϵ^*}) is bounded from above by the sum of $m = |Q|$ i.i.d. indicator random variables with expectation $\rho_{\epsilon^*} = \text{Vol}(\overline{F}_{\epsilon^*})$. Using Lemma 5.3, it is easy to show that this sum is

$$O(m\rho_{\epsilon^*} + \log n)$$

with probability $1 - n^{-C}$ where one can have the constant C arbitrarily large.

The handling of $Z(P) - Z(P')$ is more delicate, as a point from $Q = P' \setminus P$ can decrease $Z(P)$ by an arbitrary amount. For $k \geq 2$, we say that a point x is *k-wide* with respect to P if x lies outside the convex hull of P and sees k vertices of this convex hull.

Define

$$U_{k,P} = \{x \mid x \text{ is } k\text{-wide with respect to } P\}.$$

The key new ingredient is the following lemma which asserts that with high probability, $U_{k,P}$ has small volume.

Lemma 8.4. *There are positive constants c_1, c_2, c_3, c_4 such that the following holds. For any $k \geq c_3$ and $T \geq c_3 \max\{\frac{k}{c_4 n}, \rho_{\frac{k}{c_4 n}} \exp(-c_2 k)\}$ we have*

$$\mathbf{P}(\text{Vol}(U_{k,P}) \geq T) \leq \exp(-c_1 nT)$$

where P is a set of n random points. Furthermore, for any constant C there is a constant c_5 such that with probability at least $1 - n^{-C}$, there is no point in K which is $c_5 \log n$ -wide with respect to P .

The first half of the lemma is Lemma 7.1 from [11]. In order to maintain the flow of the presentation, we defer the proof of the second half to the appendix.

Let c_6 be a large positive constant. For each $c_3 \leq k \leq c_5 \log n$, define

$$T_k = \max\{c_3 \max\{\frac{k}{c_4 n}, \rho_{\frac{k}{c_4 n}} \exp(-c_2 k)\}, c_6 \log n/n\}.$$

The first statement of Lemma 8.4 implies that with probability at least

$$1 - \sum_{k=c_3}^{c_5 \log n} \exp(-c_1 c_6 \log n)$$

we have

$$\text{Vol}(U_{k,P}) \leq T_k$$

for all $c_3 \leq k \leq c_5 \log n$.

We call the random set P *perfect* if the following three conditions hold

- $\text{Vol}(U_{k,P}) \leq T_k$ for all $c_3 \leq k \leq c_5 \log n$.
- $\text{Vol}(U_{k,P}) = 0$ for any $k > c_5 \log n$.
- $\text{Vol}(K \setminus P) \leq c_7 \rho_{1/n}$.

The above argument and the second part of Lemma 8.4 and Theorem 5.1 together imply that for any given constant C , we can choose the constants c_3, c_5 and c_7 such that P is perfect with probability at least $1 - n^{-C}$, (The values of these constants do not play any role in what follows.)

Now consider a perfect set P . We are going to expose a set Q of m random points and see how it changes the number of vertices of $\text{Conv}(P)$. Let x be a random point and S be the number of vertices of $\text{Conv}(P)$ seen by x . The distribution of S , due to the fact that P is perfect, satisfies

$$\mathbf{P}(S \geq k) \leq T_k$$

for $c_3 \leq k \leq c_5 \log n$ and

$$\mathbf{P}(S \geq k) = 0$$

for $k > c_5 \log n$. Furthermore, for any $1 \leq k < c_3$,

$$\mathbf{P}(S = k) \leq \text{Vol}(K \setminus P) \leq c_7 \rho_{1/n}.$$

The difference $X = Z(P) - Z(P')$, conditioned on a perfect set P , is bounded from above by the sum of m independent copies of S . A routine Laplace transform argument gives

$$\mathbf{P}(X \geq D) = \mathbf{P}(e_0^X \geq e_0^D) \leq \mathbf{E}(e_0^X) e_0^{-D} = \mathbf{E}(e_0^S)^m e_0^{-D} \quad (12)$$

for any number $e_0 > 1$. It remains to bound $\mathbf{E}(e_0^S)$, for some special choice of e_0 . Observe that

$$\mathbf{E}(e_0^S) \leq 1 + \sum_{k=1}^{\infty} e_0^k \mathbf{P}(S = k). \quad (13)$$

The information about the distribution of S enables us to bound this from above by

$$1 + O(\rho_{1/n}) + \sum_{k=c_3}^{c_5 \log n} e_0^k T_k. \quad (14)$$

By the definition of T_k , there is a constant c_8 such that

$$\sum_{k=c_3}^{c_5 \log n} e_0^k T_k \leq c_8 \sum_{k=c_3}^{c_5 \log n} e_0^k \left(\frac{k}{c_4 n} + \rho_{\frac{k}{c_4 n}} \exp(-c_2 k) + \frac{\log n}{n} \right). \quad (15)$$

Set e_0 sufficiently close to one, we have

$$\sum_{k=c_3}^{c_5 \log n} e_0^k \frac{k}{c_4 n} \leq n^{-.9} \text{ and } \sum_{k=c_3}^{c_5 \log n} e_0^k \frac{\log n}{n} < n^{-.9}. \quad (16)$$

Again by setting e_0 close to one, we can assume that $e_0^k \exp(-c_2 k) \leq \exp(-c_2 k/2)$. Thus,

$$\sum_{k=c_3}^{c_5 \log n} e_0^k T_k = O\left(\sum_{k=c_3}^{c_5 \log n} \rho_{\frac{k}{c_4 n}} \exp(-c_2 k/2) + n^{-.9} \right).$$

Using the facts that

$$\rho_\epsilon = \Theta(\epsilon^{2/(d+1)}) \text{ and } \int_{c_3}^{\infty} x^{2/(d+1)} \exp(-c_2 x/2) \partial x \text{ converges,}$$

one can easily see that

$$\sum_{k=c_3}^{c_5 \log n} \rho_{\frac{k}{c_4 n}} \exp(-c_2 k/2) = O(n^{-2/(d+1)}) = O(\rho_{1/n}). \quad (17)$$

The error term $n^{-.9}$ is negligible. Thus we can conclude that there is a positive constant γ such that for any $1 < e_0 < 1 + \gamma$

$$\mathbf{E}(e_0^S) = 1 + O(\rho_{1/n}).$$

By (12), the probability $\mathbf{P}(X \geq D)$ is bounded from above by

$$\mathbf{E}(e_0^S)^m e_0^{-D} = \exp(O(m\rho_{1/n}) - D \log e_0) \leq \exp(c_9 m \rho_{1/n} - D \log e_0) \quad (18)$$

for some constant c_9 which depends on the choice of e_0 . Setting

$$D = \frac{1}{\log e_0} (c_9 m \rho_{1/n} + C \log n)$$

(where C is a large constant) we conclude that

$$\mathbf{P}(X \geq D) \leq n^{-C}.$$

The rest of the proof is the same as in the proof for volume. The value of D is negligible compared to σ , the standard deviation of Z . Indeed, recall that the latter is $\Omega(n^{(d-1)/2(d+1)})$ (see [7]) and $m = O(\sqrt{n \log n})$. Thus,

$$D = O(\sigma n^{-1/(d+1)} \sqrt{\log n}).$$

This, in particular, shows that we can set $\delta = 1/(d+1) + o(1)$ as claimed in the theorem. The reader is invited to fill in the details. $\square$

9. PROOF OF THEOREM 1.3 (GENERAL CASE)

In order to obtain Theorem 1.3 in full generality, we need to make a few technical modifications. Let us denote by Z_i the number of facets of dimension i of K_n (thus $Z = Z_0$). Similarly, we define S_i and X_i as the analogues of S and X , respectively. The main reason for modification is that S_i can be significantly larger than $S_0 = S$.

To start, we have the following generalization of Theorem 9.1, which we can prove by slightly modifying the proof of Theorem 9.1 from [11].

Theorem 9.1. *Let K be a smooth convex body with volume one in $\mathbb{R}^d$. For any $0 \leq i \leq d-1$, there are positive constants c_i and γ_i and β_i such that the following holds. For any $0 < \lambda < \gamma_i n^{\beta_i}$,*

$$\mathbf{P}(|Z_i - \mathbf{E}(Z_i)| \geq \sqrt{\lambda n^{(d-1)/(d+1)}}) \leq 2 \exp(-c_i \lambda).$$

We are going to use the following lemma, which is a corollary of the classical Upper Bound theorem.

Lemma 9.2. *For every $0 \leq i < d$, let $\alpha_i = \min\{i, d-i\}$. Then*

$$|S_i| = O(|S_0|^{\alpha_i}).$$

We are going to consider the following analogue of (12)

$$\mathbf{P}(X_i \geq D_i) = \mathbf{P}(e_i^{X_i} \geq e_i^{D_i}) \leq \mathbf{E}(e_i^{X_i}) e_i^{-D_i} = \mathbf{E}(e_i^{S_i})^m e_i^{-D_i} \quad (19)$$

where e_i and D_i are going to be defined. Similar to (13), we have

$$\mathbf{E}(e_i^{S_i}) \leq 1 + \sum_{k=1}^{\infty} e_i^k \mathbf{P}(S_i = k). \quad (20)$$

Using the information about the distribution of $S_0 = S$ and Lemma 9.2, we can bound the right hand side from above by

$$1 + O(\rho_{1/n}) + O\left(\sum_{k=c_3}^{c_5 \log n} e_i^{\beta k^{\alpha_i}} T_k k^{\alpha_i-1}\right) \quad (21)$$

for some constants β , c_3 and c_5 . The key difference (compared to the case $i = 0$) here is that the exponent of e_i is now a polynomial in k . Thus, we need to find some suitable values for e_i and D_i . We set

$$e_i = 1 + \frac{\epsilon_i}{\log^{\alpha_i} n}$$

for some sufficiently small positive constant ϵ_i . By the definition of T_k , there is a constant c_8 such that

$$\sum_{k=c_3}^{c_5 \log n} e_i^{\beta k^{\alpha_i}} T_k k^{\alpha_i-1} \leq c_8 \sum_{k=c_3}^{c_5 \log n} e_i^{\beta k^{\alpha_i}} \left(\frac{k}{c_4 n} + \rho_{\frac{k}{c_4 n}} \exp(-c_2 k) + \frac{\log n}{n} \right) k^{\alpha_i-1}. \quad (22)$$

Set the constant ϵ_i in the definition of $e_i = \frac{\epsilon_i}{\log^{\alpha_i} n}$ sufficiently small, we have

$$\sum_{k=c_3}^{c_5 \log n} e_i^{\beta k^{\alpha_i}} \frac{k^{\alpha_i}}{c_4 n} \leq n^{-.9} \text{ and } \sum_{k=c_3}^{c_5 \log^{\alpha_i} n} e_i^{\beta k^{\alpha_i}} k^{\alpha_i-1} \frac{\log n}{n} < n^{-.9}. \quad (23)$$

and also $e_i^{\beta k^{\alpha_i}} \exp(-c_2 k) \leq \exp(-c_2 k/2)$. Thus,

$$\sum_{k=c_3}^{c_5 n} e_i^{\beta k^{\alpha_i}} T_k k^{\alpha_i-1} = O\left(\sum_{k=c_3}^{c_5 \log n} \rho_{\frac{k}{c_4 n}} \exp(-c_2 k/2) k^{\alpha_i-1} + n^{-.9} \right).$$

Using the facts that

$$\rho_\epsilon = \Theta(\epsilon^{2/(d+1)}) \text{ and } \int_{c_3}^{\infty} x^{\alpha_i-1+2/(d+1)} \exp(-c_2 x/2) dx \text{ converges,}$$

we obtain the following analogue of (17)

$$\sum_{k=c_3}^{c_5 \log n} \rho_{\frac{k}{c_4 n}} \exp(-c_2 k/2) k^{\alpha_i-1} = O(n^{-2/(d+1)}) = O(\rho_{1/n}). \quad (24)$$

Thus we have the following analogue of (18)

$$\mathbf{P}(X_i \geq D_i) \leq \mathbf{E}(e_i^{S_i})^m e_i^{-D_i} e_i^{-D_i} = \exp(-c_9 m \rho_{1/n} - D_i \log e_i). \quad (25)$$

In order to make the right hand side at most n^{-C} , for a sufficiently large constant C , we need to set

$$D = \frac{1}{\log e_i} (c_9 m \rho_{1/n} + C \log n).$$

Recall that $e_i = 1 + \Omega(\frac{1}{\log^{\alpha_i} n})$. Thus,

$$D = O(m \rho_{1/n} + \log n) \log^{\alpha_i} n = O(\sigma_i n^{-1/(d+1)} \log^{1/2+\alpha_i} n),$$

where $\sigma_i = \Theta(n^{(d-1)/2(d+1)})$ is the standard deviation of Z_i . This concludes the proof. $\square$

10. A SUMMARY

In this section we give a brief summary about the volume and the number of facets of a random polytope in a smooth convex body. Most of these results (except those concerning expectation) have been obtained in the last three years or so, using tools from modern probability. In this theorem, we use the notation

$$M_k = \int |\text{Vol}(K_n) - \mathbf{E}(\text{Vol}(K_n))|^k.$$

For k even, M_k is the k th moment.

Volume.

Theorem 10.1. *Let K be a smooth convex body of volume one in $\mathbb{R}^d$.*

(Expectation)

- *There is a positive constant c_K such that*

$$\mathbf{E}(\text{Vol}(K_n)) = 1 - (c_K + o(1))n^{-2/(d+1)}.$$

(Variance)

- $\mathbf{Var}(\text{Vol}(K_n)) = \Theta(n^{-(d+3)/(d+1)})$.

(Higher Moments)

- *For any fixed k , $M_k(\text{Vol}(K_n)) = \Theta(n^{-k+(d-1)/(d+1)})$. For any fixed odd k , absolute value of the k th moment of $\text{Vol}(K_n)$ is*

$$O(n^{-k+(d-2+o(1))/(d+1)}) = O(M_k n^{-1/(d+1)+o(1)}).$$

(Speed of Convergence)

- *Almost surely,*

$$\lim_{n \rightarrow \infty} \left| \frac{\text{Vol}(K_n)_n}{\mathbf{E}(\text{Vol}(K_n)_n)} - 1 \right| \delta(n) n^{(d+3)/(d+1)} \ln^{-1/2} n = 0$$

for any function $\delta(n)$ tending to zero with n .

(Central Limit Theorem)

- $\text{Vol}(K_n)$ *satisfies the central limit theorem.*

(Exponential Tail)

- *There is a positive constant c such that for any*

$$0 < \lambda \leq \alpha n^{\frac{(d-1)(d+3)}{(d+1)(3d+5)}}$$

we have

$$\mathbf{P}(|\text{Vol}(K_n) - \mathbf{E}(\text{Vol}(K_n))| \geq \sqrt{\lambda \mathbf{Var}(\text{Vol}(K_n))}) \leq 2 \exp(-c\lambda) + \exp(-cn^{(d-1)/(3d+5)}).$$

Remark 10.2. The statement on the expectation is due to Bárány [1] and Schütt [10]. The statement on the variance is due to Reitzner [7, 8]. For the upper bound on the variance, Vu [11] has a different proof, which also extends to M_k for any fixed k . The lower bound for M_k follows from the central limit theorem. The estimate for odd moments follows the bound on M_k and the error term ϵ_n in Theorem 1.2. The statement concerning the central limit theorem is Theorem 1.2. The statement concerning exponential tail follows from Theorem 5.1. The statement on the speed of convergence is a corollary of this theorem.

Facets.

Theorem 10.3. *Let K be a smooth convex body of volume one in $\mathbb{R}^d$ and $0 \leq i < d$ be an integer.*

(Expectation)

- *There is a positive constant c_i such that*

$$\mathbf{E}(f_i) = (c_i + o(1))n^{(d-1)/(d+1)}.$$

(Variance)

- $\mathbf{Var}(f_i) = \Theta(n^{(d-1)/(d+1)})$.

(Higher Moments)

- *For any fixed k , $M_k(f_i) = \Theta(n^{k(d-1)/2(d+1)})$. For any fixed odd k , the absolute value of the k th moment of f_i is*

$$O(n^{(k(d-1)-2)/2(d+1)+o(1)}) = O(M_k n^{-1/(d+1)+o(1)}).$$

(Speed of Convergence)

- *Almost surely,*

$$\lim_{n \rightarrow \infty} \left| \frac{f_i}{\mathbf{E}(f_i)} - 1 \right| \delta(n) n^{-(d-1)/(d+1)} \ln^{-1/2} n = 0$$

for any function $\delta(n)$ tending to zero with n .

(Central Limit Theorem)

- *f_i satisfies the central limit theorem.*

(Exponential Tail)

- *There are positive constants c_i, γ_i, β_i such that for any*

$$0 < \lambda \leq \gamma_i n^{\beta_i}$$

we have

$$\mathbf{P}(|f_i - \mathbf{E}(f_i)| \geq \sqrt{\lambda \mathbf{Var}(f_i)}) \leq 2 \exp(-c_i \lambda).$$

Remark 10.4. The asymptotic of the expectation is from Reitzner [9]. Two special cases $i = 0$ and $i = d - 1$ were established earlier by Bárány [1] and Weieacker [12], respectively. The origin of the remaining statements is the same as in the previous remark.

11. MORE CENTRAL LIMIT THEOREMS

As the reader has possibly noticed, the key ingredients in our approach are: the results in Section 2 concerning the Poisson model, the concentration results and the bounds on the differences. Once these ingredients are available, we are ready to prove a central limit theorem.

A nice thing about the current method is that the proofs involving in these ingredients are quite robust, at least in spirit. They involve very abstract tools which have little to do with the specific model of the random polytope one deals with. Representative examples are the Baldi-Rinott theorem used in [7] and the Divide and Conquer Martingale technique used in [11]. Thus, typically we can extend the results obtained here for many other models of random polytopes (the actual argument may get complicated in some occasions). For instance, analogues of the results in Section 2 are now available for the case when the body K is a polytope, due to a recent result of Bárány and Reitzner (private communication from Bárány). Furthermore, concentration results are ready for this model as well (see [11]). Thus one can prove the corresponding central limit theorems. Details will appear elsewhere.

12. APPENDIX

Proof of the second half of Lemma 8.4. Consider a floating body $F = F_{c_0 \log n/n}$ for some constant c_0 . An observation from [11] shows that there is a collection of $m = O(n^{c_1})$ caps $C_1, \dots, C_m$, each of volume $c_2 \log n/n$, where both c_1 and c_2 are constants, such that the following holds: Let x be a point in $K \setminus F$. Then the region seen by x is contained in one of the C_i .

Now let us fix a constant C as in Lemma 8.4. Choose c_0 sufficiently large so that the probability that $F_{c_0 \log n/n} \subset K_n$ is at least $1 - n^{-C-1}$. The choice of c_0 determines c_1 and c_2 . Next, we choose c_5 sufficiently large so that the probability that each of the C_i contains at least $c_5 \log n$ points is at most n^{-C-c_1-1} .

We say that a set P of n random points is regular if $F_{c_0 \log n/n} \subset \text{Conv}(P)$ and each C_i contains less than $c_5 \log n$ points from P .

It is clear that the probability that there is no point which is $c_5 \log n$ -wide with respect to P is at least the probability that P is regular, which is bounded from below by

$$1 - n^{-C-1} - mn^{-C-c_1-1} = 1 - n^{-C-1} - n^{c_1}n^{-C-c_1-1} \geq 1 - n^{-C},$$

completing the proof. □

REFERENCES

- [1] I. Bárány, Random polytopes in smooth convex bodies, *Mathematika* **39** (1992) 81-92.
- [2] I. Bárány and L. Dalla, Few points to generate a random polytope, *Mathematika* **44** (1997), no. 2, 325–331.
- [3] I. Bárány and D. Larman, Convex bodies, economic cap coverings, random polytopes, *Mathematika* **35** (1988), no. 2, 274–291.
- [4] B. Efron, The convex hull of a random set of points, *Biometrika* **52** (1965) 331–343.
- [5] T. Hsing, On the asymptotic distribution of the area outside a random convex hull in a disk, *Ann. Appl. Probab.* **4** (1994), no. 2, 478–493.
- [6] A. Rényi and R. Sulanke, ber die konvexe Hlle von n zufllig gewhlten Punkten, *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **2**, 75–84 (1963).
- [7] M. Reitzner, Central Limit Theorems for Random Polytopes, *Probability and related fields*, to appear.
- [8] M. Reitzner, Random polytopes and the Efron-Stein Jackknife inequality, *Annals of probability*, **31** (2003), 2136–2166.
- [9] M. Reitzner, The combinatorial structure of random polytopes, *Adv in Math.* **191** (2005) 178–208.
- [10] C. Schütt, Random polytopes and affine surface area, *Math. Nachr.* **170** (1994) 227–249.
- [11] V. H. Vu, Sharp concentration of random polytopes, *GAFA*, to appear.
- [12] J. Wieacker, Einige Probleme der polyedrischen Approximation, Freiburg im Breisgau, Diplomarbeit, 1978.
- [13] W. Weil and J. Wieacker, Stochastic geometry, Handbook of convex geometry, Vol. A, B, 1391–1438, North-Holland, Amsterdam, 1993.

DEPARTMENT OF MATHEMATICS, UCSD, LA JOLLA, CA 92093-0112

E-mail address: `vanvu@ucsd.edu`